

## Photogravure

## Distribution of Urocortin 2 and Urocortin 3 in Rat Brain

Asuka Mano-Otagiri and Tamotsu Shibasaki  
Department of Physiology, Nippon Medical School




Fig. 1

Recently, urocortin (Ucn) 2 and Ucn 3 showing high sequence homologies to corticotropin-releasing factor (CRF) have been discovered as members of CRF peptide family<sup>1,3</sup>. Both peptides bind specifically to CRF<sub>2</sub> receptor. We generated polyclonal antisera against Ucn 2 and Ucn 3 to define distribution patterns of these peptides in rat central nervous system.

Many Ucn 2 neurons are present in both parvocellular and magnocellular divisions of the paraventricular nucleus (PVN) and a few Ucn 2 neurons exist in the supraoptic nucleus (SON) of the hypothalamus. Ucn 3 neurons exist in the rostral perifornical area between the fornix and the PVN, with a few cells found in the magnocellular division of the PVN. In the extra-hypothalamic region, Ucn 3 neurons are found in the dorsal division of the medial nucleus of the amygdala. These distribution patterns suggest distinct functions of Ucn 2 and Ucn 3 in the brain.

## References

1. Reyes TM, Lewis K, Perrin HM, Kunitake KS, Vaughan J, Aruas CA, Hogenesch JB, Gulyas J, Rivier J, Vale WW, Sawchenko PE: Urocortin II: A member of the corticotropin-releasing factor (CRF) neuropeptide family that is selectively bound by type 2 CRF receptors. *Proc Natl Acad Sci USA* 2001; 98: 2843-2848.
2. Lewis K, Perrin MH, Blount A, Kunitake K, Donaldson C, Vaughan J, Reyes TM, Gulyas J, Fischer W, Bilezikjian L, Rivier J, Sawchenko PE, Vale WW: Identification of urocortin III, an additional member of the corticotropin-releasing factor (CRF) family with high affinity for CRF<sub>2</sub> receptor. *Proc Natl Acad Sci USA* 2001; 98: 7570-7575.
3. Hauger RL, Grigoriadis DE, Dallman MF, Plotsky PM, Vale WW, Dautzenberg FM: International union of pharmacology. XXXVI. Current status of the nomenclature for receptors for corticotropin-releasing factor and their ligands. *Am Soc Pharm Exp Ther* 2003; 55: 21-26.

Correspondence to Asuka Mano-Otagiri, Department of Physiology, Nippon Medical School, 1-1-5 Sendagi, Bunkyo-ku, Tokyo 113-8603, Japan

E-mail: asuka@nms.ac.jp

Journal Website ( <http://www.nms.ac.jp/jnms/> )





Fig. 2

Fig. 1 Ucn 2 neurons in the hypothalamus  
 Ucn 2 neurons are mainly distributed in the parvocellular and magnocellular divisions in the PVN ( A ) A small number of Ucn 2 neurons are present in the SON ( B )  
 OX: optic chiasm, BV: blood vessel, 3V: third ventricle, PVN: paraventricular nucleus, SON: supraoptic nucleus  
 Fig. 2 Ucn 3 neurons in the brain  
 Ucn 3 neurons are mainly distributed in the rostral perifornical area between the fornix and the PVN, with a few cells found in the magnocellular division of the PVN ( A ) In the extra- hypothalamic region, Ucn 3 neurons are found in the dorsal division of the medial nucleus of the amygdala ( B )  
 Opt: optic tract, 3V: third ventricle, MEA: medial nucleus of the amygdala